

## Vutrisiran: Post-Dose Monitoring

The following information is provided in response to your unsolicited inquiry. It is intended to provide you with a review of the available scientific literature and to assist you in forming your own conclusions in order to make healthcare decisions. This document is not for further dissemination or publication without authorization.

The full Prescribing Information for AMVUTTRA® (vutrisiran) is provided [here](#). Alnylam Pharmaceuticals does not recommend the use of its products in any manner that is inconsistent with the approved Prescribing Information. This resource may contain information that is not in the approved Prescribing Information.

If you are seeking additional scientific information related to Alnylam medicines, you may visit the Alnylam US Medical Affairs website at [RNAiScience.com](https://RNAiScience.com).

### SUMMARY

- Monitoring of patients after administration of vutrisiran was not required in the HELIOS-A and HELIOS-B studies.<sup>1,2</sup>

### INDEX

[Clinical Data](#) – [Abbreviations](#) – [References](#)

### CLINICAL DATA

#### HELIOS-A Study

HELIOS-A was a phase 3, global, randomized, open-label study designed to evaluate the efficacy and safety of vutrisiran in patients with hATTR-PN. Patients were randomized (3:1) to receive either vutrisiran 25 mg every 3 months by subcutaneous injection (n=122) or patisiran 0.3 mg/kg every 3 weeks by IV infusion (as a reference group, n=42) for 18 months. This study used the placebo arm of the APOLLO study as an external control arm (n=77) for the primary endpoint and most other efficacy endpoints. The primary endpoint was the change from baseline in mNIS+7 at 9 months.<sup>3</sup>

Per the HELIOS-A study protocol, monitoring of patients after administration of vutrisiran was not required. Vutrisiran injections were to be administered by a trained healthcare professional.<sup>1</sup>

#### HELIOS-B Study

HELIOS-B was a phase 3, global, randomized, double-blind, placebo-controlled, multicenter study designed to evaluate the efficacy and safety of vutrisiran in patients with ATTR-CM, including both hATTR and wtATTR. Patients were randomized (1:1) to receive either vutrisiran 25 mg (n=326) or placebo (n=329) every 3 months by subcutaneous injection for up to 36 months. The primary endpoint was the composite endpoint of all-cause mortality and recurrent CV events (CV hospitalizations and urgent heart failure visits) at the end of the double-blind treatment period in the overall population and in the monotherapy population (patients not receiving tafamidis at baseline). After the double-blind treatment period, all eligible patients remaining on the study were allowed to receive vutrisiran in an OLE.<sup>4</sup>

Per the HELIOS-B study protocol, monitoring of patients after administration of vutrisiran was not required. Vutrisiran injections were to be administered by a trained healthcare professional.<sup>2</sup>

## ABBREVIATIONS

ATTR-CM = transthyretin amyloidosis with cardiomyopathy; CV = cardiovascular; hATTR = hereditary transthyretin amyloidosis; hATTR-PN = hereditary transthyretin amyloidosis with polyneuropathy; IV = intravenous; mNIS+7 = modified Neuropathy Impairment Score +7; OLE = open-label extension.

*Updated 05 February 2025*

## REFERENCES

1. Alnylam Pharmaceuticals. Data on file. MED-ALL-TTRSC02-2300015.
2. Protocol for: Fontana M, Berk JL, Gillmore JD, et al. Vutrisiran in patients with transthyretin amyloidosis with cardiomyopathy. *N Engl J Med*. 2025;392(1):33-44. doi:10.1056/NEJMoa2409134
3. Adams D, Tournev IL, Taylor MS, et al. Efficacy and safety of vutrisiran for patients with hereditary transthyretin-mediated amyloidosis with polyneuropathy: a randomized clinical trial. *Amyloid*. 2023;30(1):18-26. doi:10.1080/13506129.2022.2091985
4. Fontana M, Berk JL, Gillmore JD, et al. Vutrisiran in patients with transthyretin amyloidosis with cardiomyopathy. *N Engl J Med*. 2025;392(1):33-44. doi:10.1056/NEJMoa2409134