

Vutrisiran Improved Outcomes Versus Placebo in Patients with Transthyretin Amyloidosis with Cardiomyopathy and Severe Chronic Kidney Disease: Post Hoc Analysis of HELIOS-B

Farooq H. Sheikh,¹ Julien Dang,² Marianna Fontana,³ Vincent Audard,⁴ Pablo Garcia-Pavia,⁵ Michel Khouri,⁶ Antoine Jobbe-Duval,⁷ Yevgeniy Brailovsky,⁸ Hua Zheng,⁹ Satish Early,⁹ Colleen Moffitt,⁹ Ali Yilmaz¹⁰

¹MedStar Heart and Vascular Institute/Georgetown University School of Medicine, Washington, DC, USA; ²Assistance Publique des Hôpitaux de Paris, Nephrology Department, Hôpital Ambroise Paré, Boulogne-Billancourt, France; ³National Amyloidosis Centre, University College London, London, UK; ⁴Assistance Publique des Hôpitaux de Paris, Nephrology Department, Henri Mondor Hospital University, University Paris Est Créteil and National Institute of Health and Medical Research, Creteil, France; ⁵Hospital Universitario Puerta de Hierro, Madrid, Spain; ⁶Duke University School of Medicine, Durham, NC, USA; ⁷Médipôle Hôpital Mutualiste, Villeurbanne, France; ⁸Columbia University Irving Medical Center, New York, NY, USA; ⁹Amylin Pharmaceuticals, Cambridge, MA, USA; ¹⁰Klinik für Kardiologie I, Sektion für Herzbildgebung, Universitätsklinikum Münster, Münster, Germany

Disclosures



- **FHS:** Reports consultancy fees and institutional research support from Alnylam, AstraZeneca, Bridgebio, Intellia, and Pfizer
- **JD:** Reports speaking fees from Alnylam and AstraZeneca
- **MF:** Reports consultancy/advisory boards for Alexion/Caelum Biosciences, Alnylam, AstraZeneca, Attralus, Bayer, BridgeBio/Eidos, Cardior, Intellia Therapeutics, Ionis Pharmaceuticals, Janssen Pharmaceuticals, Lexeo Therapeutics, Mycardium, Novo Nordisk, Pfizer, and Prothena; research grants from Alnylam, AstraZeneca, BridgeBio, and Pfizer; salary from the British Heart Foundation Intermediate Fellowship; share options in Lexeo Therapeutics; and shares in Mycardium
- **VA:** Reports consulting fees from Addmedica, Alnylam, AstraZeneca and Vifor outside of the submitted work
- **PG-P:** Reports speaking fees from Alnylam Pharmaceuticals, AstraZeneca, Bayer, Bridgebio, Intellia, Ionis Pharmaceuticals, NovoNordisk, and Pfizer, consulting fees from Alexion, Alnylam Pharmaceuticals, AstraZeneca, ATTRalus, Bayer, Bridgebio, Intellia, Ionis Pharmaceuticals, Life Molecular Imaging, Pfizer, Neuroimmune, and NovoNordisk, and research/educational support to my institution from Alnylam Pharmaceuticals, AstraZeneca, Bayer, Bridgebio, Intellia, NovoNordisk, and Pfizer
- **MK:** Reports speaking fees from Alnylam Pharmaceuticals and BridgeBio; consulting fees from Alnylam Pharmaceuticals, BridgeBio, and Pfizer; institutional research support from Alexion, Alnylam Pharmaceuticals, AstraZeneca, ATTRalus, BridgeBio, Intellia, Ionis Pharmaceuticals, and Pfizer
- **AJ-D:** Reports speaking/consulting fees from Alnylam Pharmaceuticals, Amicus Therapeutics, AstraZeneca, Bayer, BMS, Pfizer, Bayer, and Sanofi-Genzyme
- **YB:** Reports consultancy/advisory boards for AstraZeneca and BridgeBio, and research support from Pfizer
- **HZ, SE, CM:** Employees of Alnylam Pharmaceuticals and may hold stock and/or stock options
- **AY:** Reports advisory board activities for Alexion, Alnylam Pharmaceuticals, AstraZeneca, Bayer, BridgeBio, GE, NovoNordisk, and Pfizer, and scientific research collaborations with Circle Cardiovascular Imaging and Philips

Introduction



Transthyretin Amyloidosis with Cardiomyopathy

- ATTR-CM is a rare, progressive, and fatal systemic disease caused by misfolded amyloidogenic TTR protein accumulating as fibrils in the heart¹
 - Patients with ATTR-CM can experience complications such as progressive HF and cardiac arrhythmias, leading to a poor QoL²

Renal Dysfunction in ATTR-CM

- Decline in renal function is common among patients with ATTR, with 17–46% of patients having CKD at the time of diagnosis,^{3–5} increasing to ~60% in patients with confirmed renal amyloid deposition^{6,7}
- Renal dysfunction is a risk factor for mortality and CV hospitalization in patients with ATTR, and renal failure is an independent predictor of mortality in patients with HF^{8,9}

Vutrisiran and HELIOS-B

- Vutrisiran, an RNAi therapeutic that reduces the hepatic synthesis of variant and wild-type TTR, has been evaluated with over 1500 patient-years of experience in patients with ATTR in the Phase 3 HELIOS-A and HELIOS-B studies¹⁰
- In the HELIOS-B study (NCT04153149), vutrisiran significantly reduced risk of the composite endpoint of ACM and recurrent CV events in patients with ATTR-CM, whilst preserving functional capacity and quality of life¹¹


Objective: To assess the impact of vutrisiran on renal function and the efficacy/safety of vutrisiran in patients who advanced to CKD Stage 4 or greater during the HELIOS-B double-blind period

HELIOS-B Analysis of Renal Function and Efficacy/Safety of Vutrisiran in Patients Advancing to CKD Stage 4 or Greater



HELIOS-B: A Randomized, Double-Blind Outcomes Study in Patients with ATTR-CM¹

HELIOS-B exclusion (at screening):
eGFR <30 mL/min/1.73 m²



Post hoc analyses in the overall and monotherapy (not on baseline tafamidis) populations, and the baseline tafamidis subgroup:


All HELIOS-B patients

- Proportion with eGFR decline ≥40% from baseline
- Change in eGFR from baseline over time

Patients advancing to CKD Stage ≥4 (eGFR <30 mL/min/1.73 m²) during the double-blind period

- ACM and recurrent CV events (M33–36; primary composite)
- ACM through up to M42
- Recurrent CV events (M33–36)
- Safety

Baseline KDIGO eGFR category^a in HELIOS-B patients




^aG1, G2, G3a, G3b, and G≥4 based on eGFR ≥ 90, 60–89, 45–59, 30–44, and < 30 mL/min/1.73 m², respectively.

Abbreviations: ACM, all-cause mortality; ATTR-CM, transthyretin amyloidosis with cardiomyopathy; CKD, chronic kidney disease; CV, cardiovascular; eGFR, estimated glomerular filtration rate; KDIGO, Kidney Disease: Improving Global Outcomes; M, month; OLE, open-label extension; q3M, once every 3 months; SC, subcutaneous. **Reference:** 1. Fontana et al. *N Engl J Med.* 2025;392:33–44.

Fewer Patients Experienced Worsening Renal Function with Vutrisiran versus Placebo in HELIOS-B



Patients Experiencing a $\geq 40\%$ Decline in eGFR during the Double-Blind Period




**p $\leq$ 0.01

Abbreviation: eGFR, estimated glomerular filtration rate.

Vutrisiran Treatment Was Associated with Trends Towards Improvement in eGFR versus Placebo over Time




Baseline eGFR <60 mL/min/1.73 m²



No. of Patients						
Placebo	129	121	120	116	111	96
Vutrisiran	135	125	126	119	113	105

Baseline eGFR ≥60 mL/min/1.73 m²



No. of Patients						
Placebo	199	183	187	182	169	161
Vutrisiran	191	177	178	170	168	158

The trends in eGFR over time in the monotherapy population and the baseline tafamidis subgroup were generally similar to those in the overall population

*p<0.05; **p<0.01.
Abbreviations: eGFR, estimated glomerular filtration rate; LS, least squares; SEM, standard error of the mean.

Baseline Characteristics Were Generally Balanced across Treatment Groups among Patients Advancing to CKD Stage 4 or Greater




	Overall Population		Monotherapy Population		Baseline Tafamidis Population	
	Placebo (n=32)	Vutrisiran (n=31)	Placebo (n=19)	Vutrisiran (n=18)	Placebo (n=13)	Vutrisiran (n=13)
Age in years, mean (SD)	76.4 (6.4)	76.3 (6.4)	76.1 (7.0)	76.7 (6.9)	76.8 (5.6)	75.8 (5.9)
Male, %	96.9	87.1	94.7	83.3	100.0	92.3
BMI at baseline, mean (SD) kg/m ²	28.14 (3.58)	27.66 (3.78)	27.89 (3.66)	26.92 (3.71)	28.52 (3.56)	28.67 (3.79)
eGFR, mL/min/1.73 m ² , mean (SD)	48.4 (10.0)	46.0 (13.3)	47.8 (10.2)	44.6 (13.7)	49.2 (10.2)	48.0 (13.1)
ATTR disease type, n (%)						
Hereditary	2 (6.3)	8 (25.8)	0	6 (33.3)	2 (15.4)	2 (15.4)
Wild-type	30 (93.8)	23 (74.2)	19 (100)	12 (66.7)	11 (84.6)	11 (84.6)

Vutrisiran Reduced the Risk of Composite ACM and Recurrent CV Events versus Placebo among Patients Who Advanced to CKD Stage 4 or Greater^a



53% Risk Reduction in Composite ACM and Recurrent CV Events during the Double-Blind Period in the Overall Population



In the overall population, vutrisiran reduced the risk of ACM by 57% during the double-blind period plus 6 months of the OLE, and the risk of recurrent CV events by 51% during the double-blind period

^aIncludes events occurring on or after advancement to CKD Stage ≥4.

Abbreviations: ACM, all-cause mortality; CI, confidence interval; CV, cardiovascular; HF, heart failure; HR, hazard ratio; OLE, open-label extension.

The Safety Profile of Vutrisiran in Patients Who Developed CKD Stage 4 or Greater Was Comparable with Placebo^a



n (%)	Overall Population		Monotherapy Population		Baseline Tafamidis Population	
	Placebo (n=32)	Vutrisiran (n=31)	Placebo (n=19)	Vutrisiran (n=18)	Placebo (n=13)	Vutrisiran (n=13)
≥1 AE	29 (90.6)	28 (90.3)	17 (89.5)	17 (94.4)	12 (92.3)	11 (84.6)
Related to study drug	1 (3.1)	2 (6.5)	0	1 (5.6)	1 (7.7)	1 (7.7)
≥1 serious AE	23 (71.9)	22 (71.0)	13 (68.4)	13 (72.2)	10 (76.9)	9 (69.2)
Related to study drug	0	0	0	0	0	0
≥1 severe AE	20 (62.5)	19 (61.3)	13 (68.4)	12 (66.7)	7 (53.8)	7 (53.8)
Related to study drug	0	0	0	0	0	0
≥1 AE leading to study drug interruption	1 (3.1)	2 (6.5)	0	2 (11.1)	1 (7.7)	0
Related to study drug	0	0	0	0	0	0
≥1 AE leading to study drug discontinuation	1 (3.1)	0	1 (5.3)	0	0	0
Related to study drug	0	0	0	0	0	0
≥1 AE leading to study withdrawal	1 (3.1)	0	1 (5.3)	0	0	0
Related to study drug	0	0	0	0	0	0
Death	14 (43.8)	9 (29.0)	8 (42.1)	6 (33.3)	6 (46.2)	3 (23.1)

The safety profile of vutrisiran in patients who advanced to CKD Stage ≥4 during the double-blind period was comparable with that established in the overall population during the HELIOS-B study¹

^aIncludes events occurring on or after advancement to CKD Stage ≥4.

Abbreviations: AE, adverse event; CKD, chronic kidney disease.

Reference: 1. Fontana et al. *N Engl J Med.* 2025;392:33–44.

Conclusions



Vutrisiran may slow eGFR decline and improve outcomes versus placebo in patients with ATTR-CM and advanced CKD

- Vutrisiran appeared to preserve renal function in patients with ATTR-CM, with fewer patients experiencing $\geq 40\%$ decreases in eGFR during the double-blind period of HELIOS-B versus placebo
- Consistent with results from the overall population,¹ treatment with vutrisiran reduced the risk of ACM and CV events versus placebo in patients with ATTR-CM who progressed to CKD Stage 4 or greater during the double-blind period of HELIOS-B
- The safety profile of vutrisiran in patients with impaired renal function was consistent with its established profile¹
- These data demonstrate vutrisiran is an effective and well-tolerated treatment option for patients with ATTR-CM experiencing impaired renal function

We thank the patients, their families, investigators, staff, and collaborators for their participation in HELIOS-B

This study was funded by Alnylam Pharmaceuticals. Medical writing assistance was provided by Georgina Collett of Adelphi Communications Ltd, UK, and funded by Alnylam Pharmaceuticals in accordance with Good Publication Practice guidelines. If you are seeking additional scientific information related to Alnylam therapeutics, US HCPs may visit the Alnylam US Medical Affairs website at RNAiScience.com. Non-US HCPs should contact medinfo@alnylam.com.

Abbreviations: ACM, all-cause mortality; ATTR-CM, transthyretin amyloidosis with cardiomyopathy; CKD, chronic kidney disease; CV, cardiovascular; eGFR, estimated glomerular filtration rate; HCP, healthcare professional.

Reference: 1. Fontana et al. *N Engl J Med*. 2025;392:33–44.