

Givosiran: Elevations in Blood Homocysteine

The following information is provided in response to your unsolicited inquiry. It is intended to provide you with a review of the available scientific literature and to assist you in forming your own conclusions in order to make healthcare decisions. This document is not for further dissemination or publication without authorization.

The full Prescribing Information for GIVLAARI® (givosiran) is provided [here](#). Alnylam Pharmaceuticals does not recommend the use of its products in any manner that is inconsistent with the approved Prescribing Information. This resource may contain information that is not in the approved Prescribing Information.

If you are seeking additional scientific information related to Alnylam medicines, you may visit the Alnylam US Medical Affairs website at RNAiScience.com.

SUMMARY

- Increases in blood homocysteine levels have occurred in patients receiving givosiran. In the OLE of the ENVISION study, blood homocysteine increased was reported in 15 of 93 (16%) patients treated with givosiran.¹
 - Per the ENVISION clinical study protocol, patients with increased blood homocysteine levels were recommended to receive a supplement containing vitamin B6.²
- Case reports of elevated homocysteine in patients treated with givosiran have been published in the literature and suggest management to include vitamin B6 supplementation.³⁻⁷
- Measure blood homocysteine levels prior to initiating treatment and monitor for changes during treatment with GIVLAARI. In patients with elevated blood homocysteine levels, assess folate, vitamins B12 and B6. Consider treatment with a supplement containing vitamin B6 (as monotherapy or a multivitamin preparation).¹

INDEX

[Background](#) – [Clinical Data](#) – [Global Safety Database](#) – [Label Information](#) – [Abbreviations](#) – [References](#)

BACKGROUND

In patients with AHP, elevations in blood homocysteine have been reported, and a correlation of higher levels of homocysteine is seen in those with greater disease activity. In addition, patients with AHP commonly have deficiencies of vitamins involved in homocysteine metabolism, such as MTHFR or CBS. These deficiencies may be increased in those with greater disease severity. The clinical relevance of homocysteine elevations in patients with AHP is unknown.^{4,8,9}

Current evidence supports the possible association between increases in blood homocysteine and givosiran treatment to be attributable to impaired trans-sulfuration pathway catalyzed by CBS. This is evidenced by a strong correlation and co-increase of homocysteine and methionine observed in case reports, as well as the effective reduction of blood homocysteine with supplementation of a CBS cofactor, such as vitamin B6 or heme.³⁻⁷

CLINICAL DATA

ENVISION Study

The ENVISION study was a phase 3, randomized, double-blind, placebo-controlled, multicenter study evaluating the efficacy and safety of givosiran in patients with a documented diagnosis of AHP. Enrolled patients were randomized on a 1:1 basis to receive subcutaneous injections of givosiran 2.5 mg/kg (N=48) or placebo (N=46) once a month for 6 months, followed by an optional 30-month OLE.¹⁰

In the OLE, patients (N=93) were initially assigned to givosiran 2.5 mg/kg monthly or givosiran 1.25 mg/kg monthly. A subsequent protocol amendment was enacted to increase the dose for all patients receiving the 1.25 mg/kg monthly dose to the 2.5 mg/kg monthly dose.¹⁰

Increased blood homocysteine was reported as an AE in a total of 15 patients (16%). Two patients discontinued givosiran treatment because of SAEs of increased blood homocysteine.¹⁰

Exploratory Plasma Biomarker Assessments in the ENVISION Study

A retrospective, post-hoc analysis of the ENVISION study was conducted to analyze patients' plasma homocysteine levels. Archived serum samples of patients who consented to exploratory biomarker assessment and had baseline samples (N=92) were analyzed.¹¹

Plasma homocysteine levels measured during treatment with givosiran in the 6-month double-blind and OLE periods are presented in **Table 1**.¹¹

Table 1. Plasma Homocysteine Levels (μmol/L) in the ENVISION Study.¹¹

	Visit						
	Baseline	Month 6	Month 12	Month 18	Month 24	Month 30	Month 36
Treatment Arm, n							
Placebo/Givosiran	44	44	40	18	13	8	30
Givosiran/Givosiran	48	47	46	22	20	0	32
Plasma Homocysteine Level, median (range)	17 (6-158)	85 (7-400)	83 (8-400)	98 (13-400)	47 (14-345)	44 (9-269)	22 (10-207)

Homocysteine elevations were not associated with changes in the efficacy or safety of givosiran. No correlation was found between¹¹:

- changes in homocysteine levels at baseline and changes in ALA or PBG levels from baseline and month 6 (Pearson correlation coefficient: -0.068; P=0.54)
- changes in homocysteine and average number of attacks observed during givosiran treatment in patients with and without significant homocysteine elevations (defined as either >2x ULN or >2x baseline; or >100 micromol/L for >12 months)
- AEs observed with homocysteine status during givosiran treatment.

An additional analysis was conducted analyzing the plasma samples of ENVISION study participants for CBS activity. At month 12, a statistically significant inverse correlation was observed between homocysteine levels and CBS activity (Spearman's $r = -0.58$, $P < 0.0001$) after treatment with givosiran.¹²

Four patients with elevated homocysteine levels started a daily multivitamin containing 3 mg of vitamin B6 as pyridoxine during the ENVISION study. Patients received vitamin B6 supplementation between

32 to 34 months. Plasma homocysteine levels prior to and following administration of vitamin B6 are presented in **Table 2**. An increase in CBS activity was observed from months 24 to 36 following vitamin B6 supplementation.¹²

Table 2. Plasma Homocysteine Concentrations (µM) in Four Patients in the ENVISION Study.¹³

Patient/Treatment Arm ^a	Visit				
	Baseline	Month 6	Month 12	Month 24	Month 36
Patient 1 Placebo	30.2	23.9	241.0	320.9	30.0
Patient 2 Placebo	21.7	26.2	372.2	356.4	96.4
Patient 3 Placebo	20.6	15.1	> 400.0	331.9	46.9
Patient 4 Givosiran	16.0	41.7	45.5	49.9	18.8

^aPatients in the placebo arm began givosiran treatment at month 6. Both the baseline and month 6 homocysteine measurements were taken from plasma samples collected prior to dosing.

GLOBAL SAFETY DATABASE

A cumulative post-marketing review of Alnylam Pharmaceuticals' global safety database did not identify any new safety concerns related to increases in blood homocysteine. None of the identified cases demonstrated an association of elevations in homocysteine with other AEs, such as thromboembolic events. The risk of clinical consequences of increased blood homocysteine levels is closely monitored through routine pharmacovigilance activities.¹⁴

GIVLAARI PRESCRIBING INFORMATION – RELEVANT CONTENT

The **WARNINGS AND PRECAUTIONS** section provides the following information¹:

Blood Homocysteine Increased

Increases in blood homocysteine levels have occurred in patients receiving GIVLAARI. In the ENVISION study, during the open label extension, adverse reactions of blood homocysteine increased were reported in 15 of 93 (16%) patients treated with GIVLAARI. The clinical relevance of the elevations in blood homocysteine during treatment with GIVLAARI is unknown. Measure blood homocysteine levels prior to initiating treatment and monitor for changes during treatment with GIVLAARI. In patients with elevated blood homocysteine levels, assess folate, vitamins B12 and B6. Consider treatment with a supplement containing vitamin B6 (as monotherapy or a multivitamin preparation).

ABBREVIATIONS

AE = adverse event; AHP = acute hepatic porphyria; ALA = γ-aminolevulinic acid; CBS = cystathionine-β-synthetase; CKD = chronic kidney disease; DB = double-blind; MTHFR = methylenetetrahydrofolate reductase; OLE = open-label extension; PBG = porphobilinogen; SAE = serious adverse event.

Updated 25 June 2025

REFERENCES

1. GIVLAARI (givosiran) Prescribing Information. Cambridge, MA: Alnylam Pharmaceuticals, Inc.
2. Protocol for: Balwani M, Sardh E, Ventura P, et al. Phase 3 trial of RNAi therapeutic givosiran for acute intermittent porphyria. *N Engl J Med*. 2020;382(24):2289-2301. doi:10.1056/NEJMoa1913147
3. Petrides PE, Klein M, Schuhmann E, et al. Severe homocysteinemia in two givosiran-treated porphyria patients: is free heme deficiency the culprit? *Annals of Hematology*. 2021;100(7):1685-1693. doi:10.1007/s00277-021-04547-3

4. To-Figueras J, Wijngaard R, García-Villoria J, et al. Dysregulation of homocysteine homeostasis in acute intermittent porphyria patients receiving heme arginate or givosiran. *J Inherit Metab Dis*. 2021;44(4):961-971. doi:10.1002/jimd.12391
5. Vassiliou D, Sardh E. Homocysteine elevation in givosiran treatment: Suggested ALAS1 siRNA effect on cystathionine beta-synthase. *J Intern Med*. 2021;290(4):928-930. doi:10.1111/joim.13341
6. Ricci, Marcacci, Cuoghi, Pietrangelo, Ventura. Hyperhomocysteinemia in patients with acute porphyrias: a possible effect of ALAS1 modulation by siRNA therapy and its control by vitamin supplementation. *Eur J Intern Med*. 2021;92:121-123. doi:10.1016/j.ejim.2021.06.023
7. Fontanellas A, Ávila MA, Arranz E, Enríquez de Salamanca R, Morales-Conejo M. Acute intermittent porphyria, givosiran, and homocysteine. *J Inherit Metab Dis*. 2021;44(4):790-791. doi:10.1002/jimd.12411
8. To-Figueras J, Lopez RM, Deulofeu R, Herrero C. Preliminary report: Hyperhomocysteinemia in patients with acute intermittent porphyria. *Metabolism*. 2010;59(12):1809-1810. doi:10.1016/j.metabol.2010.05.016
9. Ventura, Corradini, Di Pierro, et al. Hyperhomocysteinemia in patients with acute porphyrias: A potentially dangerous metabolic crossroad? *Eur J Intern Med*. 2020;79:101-107. doi:10.1016/j.ejim.2020.04.002
10. Kuter DJ, Bonkovsky HL, Monroy S, et al. Efficacy and safety of givosiran for acute hepatic porphyria: Final results of the randomized phase III ENVISION trial. *J Hepatol*. 2023;79(5):1150-1158. doi:10.1016/j.jhep.2023.06.013
11. Ventura P, Sardh E, Longo N, et al. Hyperhomocysteinemia in acute hepatic porphyria (AHP) and implications for treatment with givosiran. *Expert Rev Gastroenterol Hepatol*. 2022;16(9):879-894. doi:10.1080/17474124.2022.2110469
12. Keibler MA, Sridharan GV, Sweetser MT, Ticau S. Elevated homocysteine is negatively correlated with plasma cystathionine β -synthase activity in givosiran-treated patients. *JIMD Reports*. 2024. doi:10.1002/jmd2.12416
13. Supplement to: Keibler MA, Sridharan GV, Sweetser MT, Ticau S. Elevated homocysteine is negatively correlated with plasma cystathionine β -synthase activity in givosiran-treated patients. *JIMD Reports*. 2024:1-10. doi:10.1002/jmd2.12416
14. Alnylam Pharmaceuticals. Data on file. MED-ALL-AS1-2400049.