

Givosiran: Timing of Dosing

The following information is provided in response to your unsolicited inquiry. It is intended to provide you with a review of the available scientific literature and to assist you in forming your own conclusions in order to make healthcare decisions. This document is not for further dissemination or publication without authorization.

The full Prescribing Information for GIVLAARI® (givosiran) is provided [here](#). Alnylam Pharmaceuticals does not recommend the use of its products in any manner that is inconsistent with the approved Prescribing Information. This resource may contain information that is not in the approved Prescribing Information.

If you are seeking additional scientific information related to Alnylam medicines, you may visit the Alnylam US Medical Affairs website at RNAiScience.com.

SUMMARY

- The recommended dose of givosiran is 2.5 mg/kg administered via subcutaneous injection once monthly; dosing is based on actual body weight.¹
- The recommended dose and dosing frequency of givosiran is based on data from a 3-part multidose Phase 1 study that evaluated the safety, tolerability, pharmacokinetics, and pharmacodynamics of givosiran in patients with AIP.²
 - Patients with AIP who had recurrent attacks (n=17) were split into 5 cohorts and randomized to receive givosiran 2.5 mg/kg or 5 mg/kg at quarterly or monthly intervals or placebo.²
 - Patients who were treated with monthly givosiran (2.5 mg/kg or 5 mg/kg) had sustained ALA and PBG reductions as well as lower peak-to-trough fluctuations compared to patients treated with quarterly givosiran (2.5 mg/kg or 5 mg/kg).²
 - The largest reduction of mean AAR, an exploratory endpoint, was observed in patients treated with monthly givosiran 2.5 mg/kg or 5 mg/kg vs quarterly givosiran and placebo.²
 - Most AEs in the Phase 1 Study were mild or moderate, occurred at similar rates across the treatment groups, and had no clear relationship to givosiran dose. The most common AEs across all 3 parts of the study were abdominal pain, nausea, diarrhea, and nasopharyngitis.²

INDEX

[Label Information](#) – [Clinical Data](#) – [Abbreviations](#) – [References](#)

GIVLAARI PRESCRIBING INFORMATION – RELEVANT CONTENT

For relevant labeling information, please refer to the following section of the [GIVLAARI US Prescribing Information](#):¹

- DOSAGE AND ADMINISTRATION Section 2.1 Recommended Dosage

CLINICAL DATA

Study Design

The Phase 1 study of givosiran was a multicenter, randomized, placebo-controlled, 3-part study (n=23 in Parts A and B; n=17 in Part C) designed to evaluate the safety, tolerability, pharmacokinetics, and pharmacodynamics of givosiran. The clinical activity of givosiran was also explored in Part C. Parts A and B were single-blind trials of single ascending and multiple ascending dose phases, respectively, in patients with AIP who were CHEs (defined as having a urinary porphobilinogen level >4 mmol/mol creatinine at screening and no attacks in the 6 months before baseline).²

Part C was a multidose, double-blind study in which patients with AIP were followed during a run-in period of 4 to 24 weeks. These patients had to have ≥1 attack prior to randomization, in addition to ≥2 attacks within the 6 months before the run-in period; or receive scheduled hemin prophylaxis at the start of the run-in period as needed. The 17 AIP patients were divided into 5 cohorts and were randomly assigned in a 3:1 ratio to receive givosiran (2.5 or 5.0 mg/kg) at quarterly or monthly intervals or placebo for 12 weeks (13 patients were assigned to receive givosiran and 4 to receive placebo). Patients were followed for an additional 12 weeks after the last injection.²


Pharmacodynamics

In Part C, in each of the four givosiran-treated cohorts (2.5 mg/kg quarterly for two doses, 2.5 mg/kg monthly for four doses, 5.0 mg/kg quarterly for two doses, or 5.0 mg/kg monthly for four doses), reduction of *ALAS1* mRNA from baseline was seen with maximal reductions of 49 (±3%), 67 (±3%), 53 (±7%), and 74 (±6%), respectively. Reduction in *ALAS1* mRNA was accompanied by a lowering of urinary ALA and PBG with a mean maximum reduction of >90% compared to baseline observed in both monthly doses. The monthly dosing regimens led to sustained urinary ALA and PBG reductions and lower peak-to-trough fluctuations when compared to the quarterly dosing regimen. There was no observed difference in reduction of urinary ALA or PBG with the increased monthly dose regimen (2.5 mg/kg versus 5.0 mg/kg).²

Exploratory Clinical Outcomes

The mean AAR was evaluated in Part C as an exploratory endpoint and is shown in **Figure 1**. Attacks were defined as those leading to hospitalization, urgent healthcare visits, or the use of intravenous hemin at home. The mean AAR during the treatment period of Part C was 7.2 for all givosiran-treated groups, compared to 16.7 in the placebo group (a 57% difference). The mean AAR was further reduced when only the 2.5 and 5.0 mg/kg monthly givosiran dosing groups were compared to placebo (83% and 75% reduction, respectively). Though a statistical analysis was not conducted, patients who received monthly givosiran at either dose had a lower mean AAR than patients who received either quarterly givosiran dose (**Figure 1**).²

Figure 1. Mean AAR.^{2,a}




Abbreviations: AAR = annualized attack rate; SEM = standard error of the mean.

^aT bars indicate standard errors.

From Sardh et al.²

The annualized number of hemin doses was also evaluated between treatment groups in Part C and is shown in **Figure 2**. Patients treated with givosiran across all groups in Part C had a mean annualized number of hemin doses of 12.1 versus 23.4 for the placebo group (a 48% reduction). When comparing the run-in period to the treatment period, givosiran treatment reduced the annualized hemin doses by a mean of 64% across all groups as compared to a 34% mean reduction in the placebo group. Though a statistical analysis was not conducted, patients who received monthly givosiran at either dose had lower mean annualized number of hemin doses than patients who received either quarterly givosiran dose (**Figure 2**).²

Figure 2. Mean Annualized Number of Hemin Doses.^{2,a}



Abbreviations: SEM = standard error of the mean.

^aT bars indicate standard errors.

From Sardh et al.²

Safety

Most AEs in the Phase 1 Study were mild or moderate, occurred at similar rates across the treatment groups, and had no clear relationship to givosiran dosing. The most common AEs across all 3 parts of the study were abdominal pain, nausea, diarrhea, and nasopharyngitis. Injection-site reactions were reported in 6 patients who received givosiran (18%) and in no patients who received placebo.² Eight SAEs occurred in 6 patients who received givosiran and none who received placebo, including abdominal pain (2 patients) and spontaneous abortion, influenza A infection, functional gastrointestinal disorder, staphylococcal bacteremia, auditory hallucination, and fatal hemorrhagic pancreatitis (1 patient each). All SAEs were assessed as unlikely to be related to givosiran treatment by the investigating physician.^{2,3}

ABBREVIATIONS

AAR = annualized attack rate; AE = adverse event; AIP = acute intermittent porphyria; ALA = aminolevulinic acid; ALAS1 = aminolevulinic acid synthase 1; CHE = chronic high excreter; PBG = porphobilinogen; SAE = serious adverse event.

Updated 25 March 2026

REFERENCES

1. GIVLAARI (givosiran) Prescribing Information. Cambridge, MA: Alnylam Pharmaceuticals, Inc.
2. Sardh E, Harper P, Balwani M, et al. Phase 1 Trial of an RNA Interference Therapy for Acute Intermittent Porphyria. *N Engl J Med*. 2019;380(6):549-558. doi:10.1056/NEJMoa1807838
3. Supplement to: Sardh E, Harper P, Balwani M, et al. Phase 1 trial of an RNA interference therapy for acute intermittent porphyria. *N Engl J Med*. 2019;380(6):549-558. doi:10.1056/NEJMoa1807838